

Wildfire-specific Fine Particulate Matter and Risk of Hospital Admissions in Urban and Rural Counties

Jia Coco Liu,^a Ander Wilson,^b Loretta J. Mickley,^c Francesca Dominici,^b Keita Ebisu,^a Yun Wang,^b Melissa P. Sulprizio,^c Roger D. Peng,^d Xu Yue,^c Ji-Young Son,^a G. Brooke Anderson,^c and Michelle L. Bell^a

Background: The health impacts of wildfire smoke, including fine particles (PM_{2.5}), are not well understood and may differ from those of PM_{2.5} from other sources due to differences in concentrations and chemical composition.

Methods: First, for the entire Western United States (561 counties) for 2004–2009, we estimated daily PM_{2.5} concentrations directly attributable to wildfires (wildfire-specific PM_{2.5}), using a global chemical transport model. Second, we defined smoke wave as ≥ 2 consecutive days with daily wildfire-specific PM_{2.5} $> 20 \mu\text{g}/\text{m}^3$, with sensitivity analysis considering 23, 28, and $37 \mu\text{g}/\text{m}^3$. Third, we estimated the risk of cardiovascular and respiratory hospital admissions associated with smoke waves for Medicare enrollees. We used a generalized linear mixed model to estimate the relative risk of hospital admissions on smoke wave days compared with matched comparison days without wildfire smoke.

Results: We estimated that about 46 million people of all ages were exposed to at least one smoke wave during 2004 to 2009 in the Western United States. Of these, 5 million are Medicare enrollees (≥ 65 years). We found a 7.2% (95% confidence interval: 0.25%, 15%) increase in risk of respiratory admissions during smoke wave days with high wildfire-specific PM_{2.5} ($> 37 \mu\text{g}/\text{m}^3$) compared with matched non smoke wave days. We did not observe an association between smoke

wave days with wildfire-specific PM_{2.5} $\leq 37 \mu\text{g}/\text{m}^3$ and respiratory or cardiovascular admissions. Respiratory effects of wildfire-specific PM_{2.5} may be stronger than that of PM_{2.5} from other sources.

Conclusion: Short-term exposure to wildfire-specific PM_{2.5} was associated with risk of respiratory diseases in the elderly population in the Western United States during severe smoke days. See video abstract at, <http://links.lww.com/EDE/B137>.

(*Epidemiology* 2017;28: 77–85)

Wildfires are a growing concern, as climate change is anticipated to increase their frequency, intensity, and spreading speed.¹ Wildfires are known to cause substantial ecologic and economic burden, and the economic costs may be underestimated because they do not account for the potentially severe impact of air pollution from wildfires on human health.² Understanding the public health impact of wildfire smoke can inform intervention-focused policies to protect population health and promote more accurate estimates of the consequences of wildfires.³

The Western United States historically suffers from wildfires⁴ due to large areas of forests, vegetation, and relatively arid weather. The burning of biomass can dramatically increase levels of toxic air pollutants, such as fine particles (PM_{2.5}).⁵ Numerous studies have demonstrated links between all-source particulate matter (PM) measured as total mass and health outcomes, especially for respiratory and cardiovascular diseases.³ Many studies have indicated that PM_{2.5} raises more human health concerns than coarse PM because the smaller particles penetrate the respiratory system more deeply.⁶

The health effects of wildfire-emitted fine particles are not well understood. Wildfire smoke can increase ambient PM levels several times higher than that on days with no wildfire sources.³ The size of fire-generated PM tends to be small, such as fine particles (PM_{2.5}).⁷ The composition of wildfire-generated PM_{2.5} may be different from PM_{2.5} from other sources, which in turn can affect toxicity.^{8,9} Wildfires are episodic, making it especially challenging to link wildfire-specific air pollution with health.

We previously performed a literature review of the small number of studies on health impact of wildfire smoke

Submitted 4 November 2015; accepted 9 September 2016.

From the ^aSchool of Forestry and Environmental Studies, Yale University, New Haven, CT; ^bDepartment of Biostatistics, T.H. Chan School of Public Health, Harvard University, Cambridge, MA; ^cSchool of Engineering and Applied Sciences, Harvard University, Cambridge, MA; ^dDepartment of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD; and ^eDepartment of Environmental & Radiological Health Sciences, College of Veterinary Medicine & Biomedical Sciences, Colorado State University, Fort Collins, CO.

Financial supported by NIH/NIEHS R21 ES022585-01, NIH R01 ES019560, NIH R01 ES024332, EPA RD 83479801, NIH R21 ES021427, NIEHS R21 ES020152, Yale Institute of Biospheric Studies Doctoral Dissertation Improvement Grant.

This publication was developed under Assistance Agreement No. 83587101, awarded by the U.S. Environmental Protection Agency to Yale University. It has not been formally reviewed by EPA. The views expressed in this document are solely those of the authors and do not necessarily reflect those of the Agency. EPA does not endorse any products or commercial services mentioned in this publication.

The authors report no conflicts of interest.

 Supplemental digital content is available through direct URL citations in the HTML and PDF versions of this article (www.epidem.com).

Correspondence: Jia Coco Liu, Room 8B, 205 Prospect St, New Haven, CT, 06511. E-mail: coco.liu@yale.edu.

Copyright © 2016 Wolters Kluwer Health, Inc. All rights reserved.

ISSN: 1044-3983/16/2801-0077

DOI: 10.1097/EDE.0000000000000556

on community populations. We found that the results on the effects of wildfires on hospital admissions were inconsistent, especially for cardiovascular diseases, in the Western United States.³ To date, most of the literature focused on a single fire episode and small population.^{10–12} It is unknown whether the health impacts of wildfire-emitted PM_{2.5} differ from those of PM_{2.5} from other sources. As a result, research that investigates health impact from wildfires on a large geographical area and over a long time is needed.

The understanding of the health impact of wildfire-related air pollution is hindered by the challenge of estimating exposure to air pollution that can be specifically attributable to wildfires. Ambient monitors measure PM_{2.5} concentration but cannot distinguish the proportion directly attributable to fires versus other sources. The majority of current wildfire-health studies used air monitoring data, which are limited in spatial (no monitors available in rural areas) and temporal resolution (generally measure every 3–6 days) and cannot isolate wildfire-specific pollution.³

Our study aimed to address many of these challenges described above. Using a chemical transport model, we could fill in the spatial and temporal gaps of monitoring data and make source attributions of the modeled PM_{2.5}. We estimated daily 2004–2009 PM_{2.5} concentrations specifically from wildfires for 561 Western United States counties and linked them to daily numbers of Medicare admissions for respiratory and cardiovascular diseases. We applied statistical methods that have not been previously used in wildfire-health studies and estimated health impacts of wildfire-specific PM_{2.5}, incorporating rural populations into statistical analysis.

METHODS

Study Domain

The study domain is the Western United States (latitude: 31–49, longitude: –101 to –125; eFigure 1; <http://links.lww.com/EDE/B109>), where wild fires occur frequently.¹³ The study region consists of 561 counties in 16 states.

Wildfire Modeling

We employed wildfire simulations from the GEOS-Chem chemical transport model (v9-01-03) to generate daily wildfire-specific PM_{2.5} levels for 6 years (2004–2009). GEOS-Chem is a global 3D atmospheric chemistry model driven by meteorology.¹⁴ It has been used to understand the pollution impact of present-day fires^{15,16} and to predict future wildfire-specific aerosols.^{1,17} The modeling integrates meteorological data from Goddard Earth Observing System (GEOS-5) of the NASA Modeling and Assimilation Office and observed wildfire area burned based on the Global Fire Emissions Database (GFED3). GFED3 combines satellite observations of fire counts, area burned, and fuel load to produce gridded, daily maps of wildfire emissions.^{18,19}

The GEOS-Chem simulation model outputs used in this study are daily (24-hour average), gridded surface

PM_{2.5} concentrations for fire seasons (May 1–October 31) 2004–2009. The grid size is 0.5 × 0.67 degrees (approximately 50 × 75 km) latitude-by-longitude. We generated estimates under two simulations: (1) the “all-source PM_{2.5}”: total PM_{2.5} levels from all sources including wildfires; and (2) “no-fire PM_{2.5}”: PM_{2.5} from all sources except the contribution from wildfires, by performing model simulations without wildfire emissions. Non fire sources for PM_{2.5} in the West include fossil fuel combustion from transportation, industry, and power plants.^{20,21} The difference between outputs from these two simulations provides an estimate of the wildfire-specific PM_{2.5} for each day and grid cell. We defined exposure based on daily wildfire-specific PM_{2.5} estimates, which may differ from the actual locations of wildfires as smoke can travel large distances.²² This model provided exposure estimates for all study subjects in the spatial domain, including those far from monitors. The results of GEOS-Chem simulations on particulate matter have been validated against observations.^{16,23} We use ground-based or aircraft measurements, not satellite data, to validate the GEOS-Chem surface PM_{2.5}, including wildfire PM_{2.5} (eAppendix Methods 2; <http://links.lww.com/EDE/B109>).

The modeled estimates of PM_{2.5} from wildfires were spatially misaligned with health and weather data, with GEOS-Chem exposure data in a gridded form, health data at the county level, and weather data at the point level (i.e., monitor location). We converted daily grid-level wildfire-specific PM_{2.5} and all-source PM_{2.5} into daily county-level values using area-weighted averaging²⁴ (eAppendix Methods 3; <http://links.lww.com/EDE/B109>). We assumed that all persons residing in a given county have the same exposure to wildfire-specific PM_{2.5} on a given day.

Hospital Admissions Data

The hospital admission data are based on billing records 2004–2009 from the Medicare Cohort Air Pollution Study (MCAPS).²⁵ Ethical review was not required for this study. We included county-level data for all Medicare beneficiaries (US residents ≥65 years) enrolled in fee-for-service plan (70.0% of all Medicare beneficiaries) in 561 counties including rural and sparsely populated counties (eFigure 1; <http://links.lww.com/EDE/B109>). The Medicare data contain daily counts of cause-specific hospital admissions by county along with detailed information on date of admission, age category, sex, race, and daily total numbers of Medicare enrollees, representing the population at risk, in each combination of age category, sex, and race. The hospital admissions counts can include repeated admissions.

We selected emergency hospital admissions for cardiovascular (CVD) and respiratory diseases as health outcomes. A visit coded as an emergency admission might not be admitted from an emergency room/department directly but the admission was emergency (admission type is emergency not elective). Previous studies connected these disease categories

with total mass PM_{2.5}.^{25–27} The ICD-9 codes of diagnoses are in eAppendix Methods 1; <http://links.lww.com/EDE/B109>.

Air Monitoring Data and Weather Data

Daily total PM_{2.5} measurements from the monitoring data, reflecting real-world PM_{2.5} from all sources, were used to calibrate the total GEOS-Chem PM_{2.5} results (“all-source” PM_{2.5}). The air monitoring data were acquired from EPA AirData (http://aqshr1.epa.gov/aqswb/aqstmp/airdata/download_files.html#Daily). When a county had measurements from multiple monitoring sites on a given day, we averaged all monitor measurements to estimate the county’s total PM_{2.5} level on that day.

Weather information was used in statistical analysis because temperature may confound health impact of air pollution.²⁸ Daily county-level weather data, including temperature and dew point temperature, were obtained from the National Centers for Environmental Information of National Oceanic and Atmospheric Administration.

Calibration

As in other chemical transport models, the GEOS-Chem PM_{2.5} estimates were biased low during extreme events, reflecting the challenge in capturing smoke plumes on fine spatial scales.²³ To address this bias, we calibrated the daily, county-level 2004–2009 modeled total PM_{2.5} estimates ([“all-source” PM_{2.5}] in all 561 counties) with the county-level total PM_{2.5} data from air monitors, by matching the quantile functions of the two datasets. This approach scales the distribution of modeled PM_{2.5} data to more closely resemble the distribution of the monitored data.²⁹ This method maintains the ordering of PM_{2.5} in the original (modeled) data (e.g., any day above the 98th percentile of PM_{2.5} in the original modeled data is above the 98th percentile in the calibrated data). This calibration process results in empirical cumulative distribution functions for the simulated total PM_{2.5} that matches that of the observed PM_{2.5}. Hence, the overall proportion of PM_{2.5} that comes from wildfire smoke is identical in the original and calibrated data. We calibrated the daily total modeled PM_{2.5} using county-average monitoring data, calculated the proportions of total modeled PM_{2.5} contributed by modeled wildfire-specific PM_{2.5} on each day, and then multiplied the calibrated total modeled PM_{2.5} with the proportions to obtain the calibrated wildfire-specific PM_{2.5}. Results from the calibration process are shown in eTable 1 and eFigure 2 (<http://links.lww.com/EDE/B109>).

Definition of a Smoke Wave

Traditionally, the short-term effects of PM_{2.5} have been investigated by estimating the association between day-to-day variations in pollutant levels with the day-to-day variation in health outcome rates. For example, some researchers applied time-series analysis to associate daily ambient air pollution exposures with daily hospital admission rates in large multicounty studies.^{25–27} However, the frequency distribution of

wildfire-specific PM_{2.5} data differs from that of traditional ambient levels of total PM_{2.5}. Absent a wildfire smoke event, the wildfire-specific PM_{2.5} level is near zero. Among all the days with an estimated wildfire-specific PM_{2.5} levels, only 28.1% have values >1 µg/m³ but levels can reach >200 µg/m³ during the wildfire days. To estimate health effects associated with rare but extreme episodes of wildfire-specific PM_{2.5}, we introduced a new modeling approach that to our knowledge has not previously been used in the wildfire-health literature.

Specifically, we first introduce the concept of “smoke wave.” The concept of smoke wave allows us to capture periods with high concentration, sporadic, and short-lived characteristics of wildfire PM_{2.5}. We define a smoke wave as at least two consecutive days with daily calibrated wildfire-specific PM_{2.5} > 20 µg/m³ (near the 98th percentile of all county days across all 561 counties). This definition is based on daily wildfire-specific PM_{2.5} levels above a designated threshold and the daily levels in all days in a smoke wave must exceed the threshold. We conducted sensitivity analyses that varied the definition of smoke wave with respect to duration and intensity; for example, we also defined smoke wave as at least one day with daily calibrated wildfire-specific PM_{2.5} > 20 µg/m³ (“single-day smoke waves”). Among all smoke-wave days, we investigated whether health impact differs on smoke wave days with different intensity and considered intensity thresholds of 23, 28, and 37 µg/m³ corresponding to the 98.5th, 99th, and 99.5th quantile of all county days across all 561 counties, respectively. We investigated whether timing within smoke waves (during the first 2 days, 3rd to 7th day, and 8th or later day of a smoke wave) affects health risks, that is, whether the health risks on an earlier day in a smoke wave differed from those for a later day in a smoke wave. We also conducted sensitivity analysis on counties with fee-for-service enrollment ≥75% among Medicare beneficiaries.

Statistical Modeling

We conducted a matched analysis to compare the hospital admission rates (number of admissions/number of Medicare fee-for-service enrollees) on smoke-wave days (exposure) and matched non smoke-wave days (no-exposure to high wildfire-specific PM_{2.5}). We chose to conduct matched analysis because the wildfire-specific PM_{2.5} exposure is episodic and occurs infrequently (1.63% days were smoke wave days among all county days). Each smoke-wave day was matched with up to three non smoke-wave days in the same county. Smoke-wave days in counties with many smoke-wave days may be matched with fewer than three non smoke-wave days when we were not able to find three suitable no-smoke-wave days. Among the total 10,080 smoke-wave days in all counties in 6 years, 9,184 were each matched with three non smoke-wave days, 697 with two non smoke-wave days, and 199 with one non smoke-wave days. We considered non smoke-wave days to be eligible match days if they are (1) within the window of 7 calendar days before or 7 days after the smoke-wave

day but primarily in a different year (before or after the year of the smoke-wave day) and (2) separated from any other smoke-wave day by at least 2 days. Among all eligible days meeting the matching criteria for a non smoke-wave day, we selected the matched non smoke-wave days at random. By matching based on a 15-day period primarily in a different year, we accounted for larger seasonal trends such as the greater propensity for wildfires to occur during the hotter and drier months. We assessed the difference in daily temperature, daily dew point temperature, and non fire $PM_{2.5}$ for exposure (smoke wave) days and no-exposure (non smoke wave) days. All statistical analyses were conducted in Rv2.15.0.

Matching reduces the effects of confounding such as from seasonal trend.³⁰ We controlled for seasonal factors by (1) including a fixed effect of study year; (2) controlling for daily temperature; and (3) using a matched approach to ensure the same seasonality of smoke-wave days and matched non smoke-wave days. The matching approach guarantees that the smoke wave and non smoke-wave days have the same distribution across season (eTable 2; <http://links.lww.com/EDE/B109>), and hence controls by design for confounding by seasonal trends. We also conducted sensitivity analysis with the statistical model not adjusting for modeled non fire $PM_{2.5}$ levels.

We investigated the relative risk (RR) of hospital admissions on the same day as a smoke wave (lag 0). We fitted a log-linear (Poisson) mixed effects regression model separately for each disease (cardiovascular or respiratory diseases) for smoke-wave days and matched non smoke-wave days across all 561 counties (details in eAppendix Methods 4; <http://links.lww.com/EDE/B109>). Similar statistical models have been applied in previous epidemiologic studies.³¹

RESULTS

Wildfire $PM_{2.5}$ Characteristics

The frequency distribution of $PM_{2.5}$ levels from wildfire sources (calibrated) differs from that of $PM_{2.5}$ from non fire sources. Levels of wildfire-specific $PM_{2.5}$ are highly skewed, with about 72% of daily county-level calibrated wildfire-specific $PM_{2.5} < 1 \mu\text{g}/\text{m}^3$. Wildfire-specific $PM_{2.5}$ has lower mean and median, but higher extremes, compared with $PM_{2.5}$ from non fire sources (Table 1). The time-series pattern of wildfire-specific $PM_{2.5}$ is mostly zero with occasional high peaks for short periods.

Smoke Wave Characteristics

Based on our definition of a smoke wave (at least two consecutive days with wildfire-specific $PM_{2.5} > 20 \mu\text{g}/\text{m}^3$), about 66% of Western United States counties (369 of 561) experienced at least one smoke wave during the 6-year period. Among the 369 counties with at least one smoke wave, on average a county had 4.6 smoke-wave days/year (Table 2). We found that the dates and locations of smoke wave days generally matched well with MODIS records of large wildfires (eFigure 4; <http://links.lww.com/EDE/B109>).

The number of smoke-wave days experienced by counties is spatially heterogeneous. Coastal California and central Idaho had the highest frequency of smoke-wave days (>10 smoke-wave days/year; Figure 1). The average wildfire-specific $PM_{2.5}$ concentration during each smoke-wave day was lower during the first 2 days of smoke waves and gradually increased over time during a smoke wave (eFigure 3; <http://links.lww.com/EDE/B109>). The median length of a smoke wave was 3 days (ranged 2 to 58). Temperatures during smoke wave days did not differ largely based on the smoke wave day's intensity (eTable 3a; <http://links.lww.com/EDE/B109>) or smoke wave length (eTable 3b; <http://links.lww.com/EDE/B109>).

Hospital Admission Summary Statistics

The study population for the 561 counties during the study timeframe (2004–2009) includes on average about 5 million Medicare enrollees per day. This population had a total of 832,244 cardiovascular admissions and 245,926 respiratory admissions during the study time frame. Within the study time frame, 369 counties had at least one smoke wave. For these counties, there were 648,789 cardiovascular admissions and 191,095 respiratory admissions. Counties that experienced a smoke wave had, on average, lower rates of hospital admissions than counties with no smoke wave (Table 3). There are 3,844,414 people exposed to at least one smoke wave, and 1,114,513 with no-exposure to smoke waves.

Association Between Wildfire $PM_{2.5}$ and Hospital Admissions

Overall, smoke waves were not associated with increased rates of cardiovascular hospital admissions. The overall association with cardiovascular admissions on a smoke-wave day compared with a non smoke-wave day was -0.74% (95% CI: -3.1% , 1.65% ; RR = 0.99). The overall association with respiratory hospital admissions on a smoke-wave day compared with a non smoke-wave day was 2.3% (95% CI: -2.2% , 7.0% ; RR = 1.0).

Smoke-wave days with different intensity (level of wildfire $PM_{2.5}$) and the various days within the smoke waves exhibited indication of trends of different health effects. Central estimates for respiratory admissions showed an increasing trend as smoke-wave day intensity increases (Figure 2B). Smoke-wave days with intensity $>37 \mu\text{g}/\text{m}^3$ (99.5th quantile) were associated with a 7.2% increase in respiratory admissions by 7.2% (95% CI: 0.25%, 15%) compared with non smoke-wave days. Therefore, more intense smoke-wave days are estimated to have higher health impacts on respiratory diseases for the study population. This association is robust to no inclusion of a variable for non fire $PM_{2.5}$ levels in the model (results not shown). Results on single-day smoke waves and counties with fee-for-service enrollment $>75\%$ are summarized in eAppendix Results 1 and 2; <http://links.lww.com/EDE/B109>.

Central estimates for CVD admissions tend to be highest during the first 2 days of a smoke wave, and decreasing over the later days within a smoke wave (Figure 3A). Respiratory admissions exhibit an opposite trend, with higher estimate in

TABLE 1. Summary Statistics for Daily GEOS-Chem PM_{2.5} Concentrations (Calibrated) from Wildfire Sources and Non Fire Sources in 561 Western United States Counties (µg/m³) During the Wildfire Season (May 1–Oct 31), 2004–2009

	Minimum	25th Percentile	Median	Mean	75th Percentile	Maximum
PM _{2.5} from wildfires	0	0.09	0.3	2.0	1.2	242
PM _{2.5} from non fire sources	0	4.4	6.2	7.0	8.7	45.1

TABLE 2. Summary Statistics for SW (Defined as at Least Two Consecutive Days with Wildfire-specific PM_{2.5} >20 µg/m³) for the 369 Western United States Counties that Experienced SW During 2004–2009

SW Characteristics	Average (SD)	Median	Minimum	Maximum
No. SW days/yr ^a	4.6 (4.9)	2.5	0.33	26.5
No. SW events/yr ^a	1.0 (0.8)	0.83	0.17	3.8
SW intensity (µg/m ³) ^b	29.3 (6.4)	28.1	20.1	70.0
SW length (days) ^b	4.4 (4.7)	3	2	58

^aStatistics based on the 369 county-average values.

^bStatistics based on all SW-level values across all SWs in the 369 counties.
SW indicates smoke waves.




FIGURE 1. Average number of smoke wave days/year for 561 Western United States counties during 2004–2009. Hashed counties have population >75,000 in the 2010 Census.

later days of the smoke wave (Figure 3B). For each types of admission, effect estimates based on timing within a smoke wave were imprecise.

DISCUSSION


Our systematic assessment indicates an association between respiratory admissions and intense smoke-wave

days, with daily wildfire-specific PM_{2.5} levels >37 µg/m³. Single-day smoke waves have a potentially more certain positive association with respiratory admission rates, possibly due to larger sample sizes and the acute response of respiratory diseases.

To our knowledge, this is the first study to use wildfire-specific data to analyze the health impact of wildfire-specific PM_{2.5}

TABLE 3. County-level Hospital Admission per 100,000 Medicare Enrollees per Day (2004–2009)

	Minimum	25th percentile	Median	Mean	75th percentile	Maximum
561 counties						
Cardiovascular disease	1.59	8.18	11.5	12.2	15.0	43.7
Respiratory	0	1.81	3.33	3.59	4.87	17.1
369 counties with smoke waves						
Cardiovascular disease	1.59	7.87	10.7	11.2	13.7	39.7
Respiratory	0	1.63	3.07	3.25	4.52	11.7
192 counties with no smoke waves						
Cardiovascular disease	4.88	9.03	13.5	14.1	17.4	43.7
Respiratory	0	2.43	3.91	4.25	5.74	17.8

**FIGURE 2.** Associations between hospital admissions and exposure to smoke wave days (compared with non-smoke-wave days) for (A) cardiovascular disease and (B) respiratory disease, by different intensity (level of wildfire-specific $\text{PM}_{2.5}$) definitions of a smoke wave. SW indicates smoke wave.

over multiple years at a large geographical scale. Key contributions of this study include (1) estimation of exposure to $\text{PM}_{2.5}$ specifically from wildfires; (2) ability to estimate exposure to wildfire $\text{PM}_{2.5}$ every county with and without air monitors, therefore expanding the study populations to include persons that live far from $\text{PM}_{2.5}$ monitoring stations; and (3) application of statistical models that estimate percent increases in hospital admission by matching smoke-wave days to non-smoke-wave days.

Although previous literature on the association between wildfire smoke and health is limited, several studies have made important contributions. The majority of such studies

used air monitor measurements, which cannot identify pollution specifically from wildfires with current technology, and studied a single wildfire episode and one or a small number of communities.³ A few studies compared air pollution exposure (from all sources) during wildfires to the periods or locations with no fire.^{11,32,33} Our study results for respiratory diseases are consistent with those found in most of the previous literatures,^{34,35} in which wildfire smoke was found to be associated with respiratory diseases. Association between wildfire smoke and cardiovascular morbidities was found in five US studies that each examined a single local wildfire episode,³ but our




FIGURE 3. Associations between hospital admissions and exposure to SW days (compared with non SW days) for (A) cardiovascular disease and (B) respiratory disease, by timing of the days within a SW. SW indicates smoke wave.

multistate, multiyear study did not provide evidence for such association.

Previous studies have demonstrated that the chemical composition of PM_{2.5}, which is related to source, can result in different effect estimates for human health.^{9,36,37} Thus, estimates from wildfire PM_{2.5} may differ from those from PM_{2.5} from other sources, such as transportation or industry. Earlier studies examined the association between risk of hospital admission and levels of PM_{2.5} from all sources (i.e., PM_{2.5} total mass; e.g., change of risk of hospital admission for Medicare enrollees per 10 µg/m³ increase in PM_{2.5} in the Western United States^{25,26,38}). As we compared the health risk among smoke-wave days with that of non smoke-wave days, rather than by a specific increment of PM_{2.5}, direct comparisons of results is challenging. Furthermore, these studies focused on urban counties with high populations, whereas our study included rural populations in the analysis as well. Still, a general comparison can give some indication of whether PM_{2.5} from wildfire smoke is more or less harmful than PM_{2.5} total mass.

For Medicare cardiovascular admissions, one study estimated an increased risk of 0.53% (95% posterior interval: 0.00%, 1.05%) per 10 µg/m³ PM_{2.5} total mass (from all sources) for 25 urban counties in the Southwest US, and 0.74% (-1.74%, 3.29%) for nine urban counties in the Northwest.²⁶ Our results did not indicate an association between wildfire PM_{2.5} and risk of cardiovascular admissions.

For respiratory hospital admissions, we estimated an increase of 7.2% (0.25%, 15%) comparing smoke-wave days with wildfire-specific PM_{2.5} > 37 µg/m³ to non smoke-wave

days with wildfire-specific PM_{2.5} ≤ 20 µg/m³, which corresponds to an average difference of 29.6 µg/m³ in those two groups of days. The earlier study identified associations between PM_{2.5} total mass and respiratory admissions for the Medicare population in the Southwest at lag 2 days at 0.94% (0.22%, 1.67%) per 10 µg/m³,²⁶ which corresponds to an increased risk of 2.8% (0.64, 5.0%) per 29.6 µg/m³. Therefore, our estimates of respiratory admissions risks indicate that wildfire-specific PM_{2.5} from intense smoke waves are associated with more harm than PM_{2.5} from other sources for the elderly in the Western United States. Further research is needed to investigate the relative toxicity of PM_{2.5} from wildfire smoke with that of other sources.

Our approaches for assessing pollutant exposure and estimating health risks address key challenges in studying the health impact of wildfire-specific pollutant. The GEOS-Chem model provided a new approach to distinguish wildfire-specific PM_{2.5} from PM_{2.5} from other sources. The fire scheme in the simulation can explain up to 60% of the observed variance of area burned in the Western United States, and is ecosystem dependent.¹⁷ This method also improves the spatial and temporal resolution of exposure estimates for air pollution. Unlike air monitoring data that generally measure PM_{2.5} concentrations every 3–6 days in urban areas, GEOS-Chem estimates concentrations for every day and covers the entire study area. Our smoke-wave methods provide an approach suitable for the study of highly-skewed air pollution data and enable identification and investigation of pollution episodes with high source-specific pollutant concentrations. Matched analysis can reduce the confounding effect of seasonality

and county-specific effects. These methods can be applied to future studies investigating other pollution events and populations.

Limitations of our study include potential spatial misalignment between the exposure estimates (gridded estimates) and health data (county). Our study population was restricted to Medicare fee-for-service enrollees, a sample of elderly persons. Our smoke wave approach does not fully capture the dose–response relationship, cause-specific health outcomes, etc., that could be investigated in future studies. The GFED emissions applied to GEOS-Chem contribute uncertainty to the modeled estimates of wildfires-specific $PM_{2.5}$. The GFED3 data may underestimate fire contributions to background $PM_{2.5}$ because of the omission of small fires³⁹ and the biases in the modeled fuel consumption. GFED3 relies on satellite observations of active fire counts and area burned, and may have difficulty discerning such phenomena, especially on cloudy days.⁴⁰ Another limitation arises as EPA monitors generally measure $PM_{2.5}$ values every 3–6 days and are located in populated areas. Given a large number of days with monitoring measurements for calibration, we assumed that the systematic sampling of EPA monitors generate measurements with mean and SD representing the full-time series of real-world $PM_{2.5}$. While it would be ideal to have the full continuous measure, we believe that calibration using this discrete sample of the continuous measure is the best possible alternative in using the available data. While our exposure estimates are advances over methods that do not isolate the air pollution from wildfires specifically, additional study could address these limitations. We choose not to a priori identify lags in this study as little is known about how wildfire-specific $PM_{2.5}$ affects human health. Most of the wildfire-health literature to date has investigated effects of lag 0 or short lags (<5 days).³ Future studies can explore the lagged effect of wildfire-specific air pollution.

Our findings indicate that wildfires are associated with increased risk of admissions for respiratory diseases for the elderly population during severe wildfire episodes. As climate change is anticipated to increase the frequency and intensity of wildfires,¹ the health burden from wildfire-specific pollutants may increase in the future. With improvement of atmospheric modeling, future studies can estimate daily wildfire-specific $PM_{2.5}$ at a finer spatial resolution. Future studies can also investigate vulnerability to wildfire smoke, health impact of different species of wildfire-specific $PM_{2.5}$, the economic consequence of the health burden from wildfire smoke, combined effect of wildfire smoke and other air pollutants, and estimated health burden in the future under climate change.

REFERENCES

1. Spracklen DV, Mickley LJ, Logan JA, et al. Impacts of climate change from 2000 to 2050 on wildfire activity and carbonaceous aerosol concentrations in the western United States. *J Geophys Res Atmos*. 2009;114. Issue D20301.
2. Kochi I, Donovan GH, Champ PA, Loomis JB. The economic cost of adverse health effects from wildfire-smoke exposure: a review. *Int J Wildland Fire*. 2010;19:803–817.
3. Liu JC, Pereira G, Uhl SA, Bravo MA, Bell ML. A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environ Res*. 2015;136:120–132.
4. Dennison PE, Brewer SC, Arnold JD, Moritz MA. Large wildfire trends in the western United States, 1984–2011. *Geophys Res Lett*. 2014;41:2928–2933.
5. Christopher JP et al.; Interagency Working Group on Climate Change and Health. In: A human health perspective on climate change: a report outlining the research needs on the human health effects of climate change. *J Current Issues Global*. 2013: 621.
6. Kim KH, Kabir E, Kabir S. A review on the human health impact of airborne particulate matter. *Environ Int*. 2015;74:136–143.
7. Janhall S, Andreae MO, Poschl U. Biomass burning aerosol emissions from vegetation fires: particle number and mass emission factors and size distributions. *Atmos Chem Phys*. 2010;10:1427–1439.
8. Bell ML, Ebisu K. Environmental inequality in exposures to airborne particulate matter components in the United States. *Environ Health Perspect*. 2012;120:1699–1704.
9. Peng RD, Bell ML, Geyh AS, et al. Emergency admissions for cardiovascular and respiratory diseases and the chemical composition of fine particle air pollution. *Environ Health Perspect*. 2009;117:957–963.
10. Delfino RJ, Brummel S, Wu J, et al. The relationship of respiratory and cardiovascular hospital admissions to the southern California wildfires of 2003. *Occup Environ Med*. 2009;66:189–197.
11. Moore D, Copes R, Fisk R, Joy R, Chan K, Brauer M. Population health effects of air quality changes due to forest fires in British Columbia in 2003: estimates from physician-visit billing data. *Can J Public Health*. 2006;97:105–108.
12. Morgan G, Sheppard V, Khalaj B, et al. Effects of bushfire smoke on daily mortality and hospital admissions in Sydney, Australia. *Epidemiology*. 2010;21:47–55.
13. Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW. Warming and earlier spring increase western U.S. forest wildfire activity. *Science*. 2006;313:940–943.
14. Bey I, Jacob DJ, Yantosca RM, et al. Global modeling of tropospheric chemistry with assimilated meteorology: model description and evaluation. *J Geophys Res Atmos*. 2001;106:23073–23095.
15. Park RJ, Jacob DJ, Logan JA. Fire and biofuel contributions to annual mean aerosol mass concentrations in the United States. *Atmos Environ*. 2007;41:7389–7400.
16. Spracklen DV, Logan JA, Mickley LJ, et al. Wildfires drive interannual variability of organic carbon aerosol in the western US in summer. *Geophys Res Lett*. 2007;34.
17. Yue X, Mickley LJ, Logan JA, Kaplan JO. Ensemble projections of wildfire activity and carbonaceous aerosol concentrations over the western United States in the mid-21st century. *Atmos Environ (1994)*. 2013;77:767–780.
18. Mu M, Randerson JT, van der Werf GR, et al. Daily and 3-hourly variability in global fire emissions and consequences for atmospheric model predictions of carbon monoxide. *J Geophys Res Atmos*. 2011;116.
19. van der Werf GR, Randerson JT, Giglio L, et al. Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmos Chem Phys*. 2010;10:11707–11735.
20. Chow JC, Watson JG. Review of $PM_{2.5}$ and PM_{10} apportionment for fossil fuel combustion and other sources by the chemical mass balance receptor model. *Energy Fuels*. 2002;16:222–260.
21. Park RJ, Jacob DJ, Field BD, Yantosca RM, Chin M. Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy. *J Geophys Res Atmos*. 2004;109.
22. Sapkota A, Symons JM, Kleissl J, et al. Impact of the 2002 Canadian forest fires on particulate matter air quality in Baltimore city. *Environ Sci Technol*. 2005;39:24–32.
23. Zhang L, Jacob DJ, Yue X, Downey NV, Wood DA, Blewitt D. Sources contributing to background surface ozone in the US Intermountain West. *Atmos Chem Phys*. 2014;14:5295–5309.

24. Bravo MA, Fuentes M, Zhang Y, Burr MJ, Bell ML. Comparison of exposure estimation methods for air pollutants: ambient monitoring data and regional air quality simulation. *Environ Res.* 2012;116:1–10.
25. Dominici F, Peng RD, Bell ML, et al. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA.* 2006;295:1127–1134.
26. Bell ML, Ebisu K, Peng RD, et al. Seasonal and regional short-term effects of fine particles on hospital admissions in 202 US counties, 1999–2005. *Am J Epidemiol.* 2008;168:1301–1310.
27. Peng RD, Chang HH, Bell ML, et al. Coarse particulate matter air pollution and hospital admissions for cardiovascular and respiratory diseases among Medicare patients. *JAMA.* 2008;299:2172–2179.
28. Samet J, Zeger S, Kelsall J, Xu J, Kalkstein L. Does weather confound or modify the association of particulate air pollution with mortality? An analysis of the Philadelphia data, 1973–1980. *Environ Res.* 1998;77:9–19.
29. Maraun D. Bias correction, quantile mapping, and downscaling: revisiting the inflation issue. *J Climate.* 2013;26:2137–2143.
30. Rubin DB. The use of matched sampling and regression adjustment to remove bias in observational studies. *Matched Sampling Causal Effects.* 2006:81–98.
31. Bobb JF, Obermeyer Z, Wang Y, Dominici F. Cause-specific risk of hospital admission related to extreme heat in older adults. *JAMA.* 2014;312:2659–2667.
32. Duclos P, Sanderson LM, Lipsett M. The 1987 forest fire disaster in California: assessment of emergency room visits. *Arch Environ Health.* 1990;45:53–58.
33. Frankenberger E, McKee D, Thomas D. Health consequences of forest fires in Indonesia. *Demography.* 2005;42:109–129.
34. Lee TS, Falter K, Meyer P, Mott J, Gwynn C. Risk factors associated with clinic visits during the 1999 forest fires near the Hoopa Valley Indian Reservation, California, USA. *Int J Environ Health Res.* 2009;19:315–327.
35. Rappold AG, Stone SL, Cascio WE, et al. Peat bog wildfire smoke exposure in rural North Carolina is associated with cardiopulmonary emergency department visits assessed through syndromic surveillance. *Environ Health Perspect.* 2011;119:1415–1420.
36. Bell ML, Ebisu K, Peng RD, Samet JM, Dominici F. Hospital admissions and chemical composition of fine particle air pollution. *Am J Respir Crit Care Med.* 2009;179:1115–1120.
37. Zanobetti A, Franklin M, Koutrakis P, Schwartz J. Fine particulate air pollution and its components in association with cause-specific emergency admissions. *Environ Health.* 2009;8:58.
38. Bell ML, Son JY, Peng RD, Wang Y, Dominici F. Ambient PM_{2.5} and risk of hospital admissions: do risks differ for men and women? *Epidemiology.* 2015;26:575–579.
39. Randerson JT, Chen Y, van der Werf GR, Rogers BM, Morton DC. Global burned area and biomass burning emissions from small fires. *J Geophys Res Biogeosci.* 2012;117.
40. Giglio L, Randerson JT, van der Werf GR. Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4). *J Geophys Res Biogeosci.* 2013;118:317–328.